

Current diagnosis and treatment in central serous chorioretinopathy

 Tuğçe Horozoğlu Ceran¹,  Mehmet Çıtırık²

¹Department of Ophthalmology, Ankara Dışkapı Yıldırım Beyazıt Training and Research Hospital, University of Health Sciences, Ankara, Turkey

²Department of Ophthalmology, Ankara Etlik City Hospital, University of Health Sciences, Ankara, Turkey

Cite this article: Horozoğlu Ceran T, Çıtırık M. Current diagnosis and treatment in central serous chorioretinopathy. *J Compr Surg.* 2023;1(2):40-44.

Corresponding Author: Tuğçe Horozoğlu Ceran, tugcehorozoglu00@gmail.com

Received: 10/04/2023

Accepted: 19/04/2023

Published: 28.05.2023

ABSTRACT

Central serous chorioretinopathy (CSCR) diagnosis and follow-up are performed with multimodal imaging methods. Fundus fluorescein angiography, optical coherence tomography (OCT), fundus autofluorescence, and indocyanine green angiography can be used in the diagnosis of patients with CSCR. OCT stands out as the gold standard for imaging, diagnosis, and follow-up. Although there is no gold standard method in treatment.

Keywords: Central serous chorioretinopathy, diagnosis, multimodal imaging, treatment modalities

INTRODUCTION

Central serous chorioretinopathy (CSCR) is a clinical condition characterized by idiopathic, well-demarcated serous detachment of the neurosensorial retina (NSR). Although serous detachment in CSCR occurs as a result of impaired barrier and pump functions of the retinal pigment epithelium (RPE), the main pathology is in the choriocapillaris.

CSCR is recognized as a pachychoroid spectrum disease. Pachychoroid spectrum diseases include focal or diffuse choroidal thickening. Thinning of the choriocapillaris and thickening of the outer choroidal layers are observed. Because of choroidal hyperpermeability, subretinal fluid can often accompany diseases in this spectrum. In addition to CSCR, pachychoroid pigment epitheliopathy, pachychoroid neovascuopathy, polypoidal choroidal vasculopathy/aneurysmal type 1 neovascularization, peripapillary pachychoroid syndrome, and focal choroidal excavation are among the diseases included in this spectrum.

CSCR can be multifocal, often involving both eyes asymmetrically. In some patients, CSCR extends to the peripheral retina, while in others it may even present as subtotal bullous retinal detachment. Thinning of the foveal contour, cystoid degeneration, and damage to the photoreceptor layer cause loss of visual function.

Although 90% of the subretinal fluid is resorbed spontaneously in 3-4 months in CSCR, the improvement in visual function may take up to 1 year. Mild to moderate metamorphopsia, indistinct scotomas, decreased contrast sensitivity and impaired color vision are among the frequently ongoing complaints.

CSCR is a clinical diagnosis. In the differential diagnosis of CSCR, polypoidal choroidal vasculopathy (PCV), CNV

(mostly Type 1), cavitory optic disc anomalies (if with serous detachment), choroidal hemangioma (if with focal choroidal thickness increase and serous detachment), dome-shaped macula (anterior protrusion of the macula with concomitant posterior staphyloma – often punctuated pooling on fluorescein angiography and increased subfoveal choroidal thickness) and Best disease (if there are splits in the RPE and elevation in the RPE) comes first.

While there is no need to request additional examination in patients with CSCR with demographic characteristics such as type A personality, age, and gender; advanced techniques such as fundus fluorescein angiography (FFA), optical coherence tomography (OCT), fundus autofluorescence (FAF), indocyanine green angiography (ICGA) can be used to support the diagnosis in patients with more severe CSCR.

IMAGING

Today, CSCR diagnosis and follow-up are performed with multimodal imaging methods. Because it is a fast and practical imaging method, OCT has often been the first imaging method. FAF is very useful for viewing RPE alterations. FFA indicates whether there is a leak or not. All imaging methods have different advantages. Therefore, multimodal imaging is important for CSCR.

FFA

The most important and most common finding in FFA is the 'expanding point' under serous retinal detachment. RPE leak occurs in 95% of CSCR cases. This leak point is 90% away from the fovea and is most common in the superonasal quadrant. In chronic CSCR cases, there is more than one

RPE leakage point.¹ Other most important characteristic FFA findings identified for CSCR are smokestack (the most classic), inkblot, and mushroom patterns.^{2,3}

OCT

The most important advantage of OCT as an imaging method is that it is non-invasive and fast. Recently, as a result of increasing the depth of imaging with OCT with EDI (enhanced depth imaging)-OCT and SS (swept source)-OCT, imaging of choroidal vessels and choroidal thickness can be performed. As choroidal changes, thickened subfoveal choroid, dilated choroidal vessels, thinned inner choroidal layer or focal choroidal excavation can be seen in CSCR.^{4,5} Choroidal thickness increased in both the affected eye and fellow eye.⁴ This increase in choroidal thickness strengthens the hypothesis that the increased hydrostatic pressure originates from the choroid.⁶ Choroidal thickness is affected by axial length and age. Choroidal thickness values above 395 micrometers are called the thick choroid (pachychoroid).⁷ Choroidal thickening results from focal or diffuse dilatation of choroidal vessels. On OCT imaging, PED, microchips, fissures, hypertrophy, or atrophy can be detected in RPE.^{4,8,9} RPE elevation occurs in the area of dilated vessels. RPE protrusion, PED, subretinal fibrinous exudate, dipping in the outer retinal layer, and elongation in the photoreceptor outer layers may occur at the leakage site.^{10,11} The area where the tiny RPE defect is seen on the OCT matches the leakage point in the FFA.¹² The elongation in the outer layers of the photoreceptor initially causes hypo-autofluorescence by suppressing the RPE, and then, with the development of atrophy (window defect), it leads to a hyper-autofluorescence image. PED appearance in OCT can be in different shapes such as dome-shaped, double dome-shaped, flat, protruding or ruptured. Intraretinal hyperreflective deposits, distortion in the ellipsoid zone, decrease in foveal thickness and cystoid macular degeneration are other findings.¹³⁻¹⁵ Focal choroidal excavation, PED, and RPE atrophy are more common in chronic CSCR.^{4,5,9} The amount of subretinal hyperreflective deposit increases with the duration of symptoms and is associated with a lower final BCVA.¹⁶ These deposits may also appear in the subretinal, intraretinal, or inner choroid.

FAF

On FAF imaging, dot-like precipitates in the outer retinal layers and subretinal yellow material are seen in CSCR.¹⁷ FAF is an imaging technique that is very useful in distinguishing acute, chronic, and recurrent cases. In acute CSCR, there is hyper-fluorescence in the detached NSR area. In chronic CSCR cases, both hyper- and hypo-fluorescent areas are seen. Areas with decreased autofluorescence do not benefit from laser treatment.¹⁸

Today, the use of FAF together with OCT is frequently preferred in clinics for the diagnosis and follow-up of CSCR.

ICGA

One of the best ways to show problems with choroidal circulation is with ICGA. Vascular abnormalities in the choroidal layer can be seen with ICGA. A delay in the filling is observed in ICGA in CSCR eyes.¹⁹ ICGA leaks from dilated choroidal vessels (congested veins).⁴ In the early phase ICGA, hypo-fluorescence is observed due to the decrease in the filling in the choriocapillaris. In mid-phase ICGA, dilated large choroidal veins are observed.

This image is due to an atrophic RPE or PED. Atrophic areas are observed as a faintly hyperfluorescent focus due to increased permeability in choroidal vessels. In the late phase, previously hyperfluorescent areas may become permanent hyperfluorescent spots or these areas may be surrounded and covered by a central displacement of another hyperfluorescent focus. There is thinning and atrophy of the choriocapillaris due to the compression effect of the dilated choroidal vessels.⁴ Hyperfluorescent areas on ICGA have non-perfused choriocapillaris.²⁰ Laser Doppler flowmetry has also shown that the blood flow in the choriocapillaris is slow.²¹

CURRENT TREATMENTS

In the treatment, the presence of risk factors that may cause CSCR should be evaluated first. Risk factors should be eliminated if possible. With the discontinuation of a used steroid, CSCR regresses by 90%.²² In the absence of risk factors that can be intervened or eliminated, the first step in the treatment of acute CSCR is follow-up because it resolves within 3-4 months.²³ There is no consensus on when to start treating a CSCR attack. Treatment is initiated in cases of persistent macular NSR detachment, which usually lasts longer than 4 months, severely reduced visual acuity, recurrence, CSCR attack causing low final BCVA in the fellow eye, and in cases where the patient is from the occupational group that needs rapid improvement of vision.²⁴

Focal Laser Photocoagulation

Laser applications at green (514 nm) and yellow (580 nm) wavelengths close the leakage point. This closure effect is provided by the induction of RPE cells and fluid efflux, as well as thermal damage.^{25,26} The effect of photocoagulation is at the RPE level and has no effect on the choroid.²⁷ After the damage created on the RPE, the surrounding RPE cells also undertake the task in that region.²⁸ Laser photocoagulation therapy does not contribute to the final BCVA and may rarely cause paracentral scotoma and iatrogenic CNV.²⁹ Therefore, it should be preferred in extrafoveal leaks.³⁰ While there are studies indicating that laser photocoagulation has no effect on recurrences,²⁹ there are studies showing that it reduces recurrences.³¹ According to the previous study, the recovery period is shortened by 2 months with photocoagulation.³¹

If the leakage point is detected, laser photocoagulation still appears to be a good treatment alternative in cases 500 microns away from the fovea and lasting longer than 4 months.

Photodynamic Therapy

Verteporfin (Visudyne, Novartis, Switzerland) is an agent that causes vessel occlusion by causing endothelial vascular damage. With the stimulation of light with a wavelength of 693 nm and oxygen, free radicals are formed in the vascular network and cause vascular damage. With this mechanism, hyperpermeable vessels in the choriocapillaris are taken under control.³² In the follow-ups, it was observed that there was a decrease in vascular permeability and even a decrease in choroidal thickness.^{33,34} In the Photodynamic Therapy (PDT) protocol applied for CSCR, application with 25 J/cm² (lower fluence, ½ fluence) setting is preferred.³⁵ Low fluence application causes fewer side effects with a good effect in CSCR treatment.³⁵ Half-dose PDT (3 mg/m²) can also be preferred to reduce side effects with a good effect.³³ 1/3-dose

PDT application is not as effective as half-dose PDT.³⁶ There is no definitive data on whether half-dose PDT or half-fluence PDT is superior. No long-term studies are showing the results and safety of PDT.

PDT results in a reduction in serous NSR detachment and symptoms, and an increase in BCVA.³⁷ Success is achieved when PDT is performed after the determination of the hyperfluorescent point after ICGA.^{38,39} PDT performed without showing the presence of hyperfluorescent spots with ICG does not have much benefit.³⁹ Complications such as a change in RPE, choriocapillaris hypoperfusion, and development of CNV may occur as a result of PDT.³⁹

Although there is a rapid regression in serous NSR detachment with low-fluence PDT at 3-month follow-up comparing PDT to focal laser photocoagulation, there is no difference in anatomical and functional success in the end.⁴⁰

Micropulse (Subliminal) Laser Photocoagulation

Subthreshold micropulse laser is also used in the treatment of CSCR.⁴¹ Induction of RPE cells is provided by giving short-term multiple pulses to the RPE cells. Higher wavelengths have a better effect on the choroid and do not damage the inner retinal layers.⁴² In some studies, subthreshold diode micropulse laser has been shown to give better results than green argon laser (faster recovery and better contrast sensitivity).

The micropulse diode laser provides an effect like PDT, but it acts more slowly.^{43,44} Iatrogenic side effects are very rare.^{43,44} However, there is no data on how long the treatment should continue.

Sub-Threshold Non-Damaging Retinal Laser Photocoagulation

The non-damaging retinal laser therapy (NRT) was recently defined. The non-damaging hyperthermia was demonstrated in mice by observing the expression of heat shock protein. Based on those reported results; the titration protocol has been developed by adjusting laser power and duration.⁴⁵ This protocol is called Endpoint Management (EpM) and is related to the minimal tissue effects in visible titration. The EpM laser therapy uses an Arrhenius integral algorithm to control laser power and pulse duration, optimizing the therapeutic effect of the laser at sub-visible levels. The laser power is titrated to create a barely visible lesion at a pulse duration of 15 or 20 milliseconds, and EpM using 15 milliseconds is called NRT.

İlhan et al.⁴⁶ treated 23 eyes of 23 treatment-naïve chronic CSCR patients with NRT. The irradiation of 577 nm yellow light was conducted on the serous detachment area after switching over to the NRT algorithm. At the 2nd-month follow-up visit after NRT, complete resorption of subretinal fluid was observed in 78.3% of eyes and incomplete resorption in 21.7% of eyes.⁴⁶ Worse values of BCVA and CMT before NRT were found to have an increased risk for incomplete resorption.⁴⁶ They concluded that significant functional and anatomical improvements can be observed in the early period after NRT in patients with chronic CSCR.⁴⁶ They emphasized that patients having worse baseline BCVA and CMT have an increased risk for incomplete resorption.⁴⁶

Selective Retinal Therapy

It acts only on intracellular melanosomes in RPE cells.⁴⁷ Since NSR is not affected, it does not cause scotoma. It has

been observed to be successful in the treatment of juxta foveal lesions.^{47,48}

Anti-VEGF

Anti-VEGF therapy has a reducing effect on choroidal permeability.⁴⁹ However, there is no increased VEGF level in CSCR.⁵⁰ Therefore, anti-VEGF should be used only in cases where CSCR is associated with CNV. With the use of bevacizumab as an anti-VEGF agent, visual acuity increases, leakage decreases, and the amount of subretinal fluid decreases.⁵¹ However, the superiority of bevacizumab treatment over other treatment methods has not been demonstrated.⁴⁹ The benefit of anti-VEGF therapy is evident in cases where CSCR is associated with CNV.⁵² While similar benefits are seen in the treatment of chronic CSCR, cases unresponsive to treatment have also been observed.⁵³ Some studies stated that anti-VEGF injection did not change the outcome of BCVA in acute CSCR.⁵⁴ Subretinal deposits are thought to be the reason for this unresponsiveness, preventing the reabsorption of fluid.⁵⁵

Mineralocorticoid Receptor Antagonist

Aldosterone and cortisol bind to mineralocorticoid receptors (MR). MR blockade is beneficial in the treatment of endothelial dysfunction, atherosclerosis, hypertension, heart failure, and chronic renal failure. In the treatment of retinopathy of prematurity, MR antagonists have an anti-angiogenic effect.⁵⁶

Spirolactone is a competitive MR antagonist drug. Spirolactone is a prodrug molecule and is converted to active metabolites in the liver. It was first used in the treatment of hypertension. Due to its binding to progesterone receptors, it causes gynecomastia, decreased libido, erectile dysfunction, and menstrual irregularities.

Eplerenone has fewer MR antagonist effects than spironolactone and has fewer hormonal side effects. Both molecules cause electrolyte imbalances and hyperkalemia. However, this side effect usually occurs in the presence of renal dysfunction.

It was observed that the foveal subretinal fluid decreased in patients treated with spironolactone and eplerenone.^{57,58}

Glucocorticoids play a role in the pathogenesis of CSCR. Therefore, anti-glucocorticoid therapy is among the treatment options for CSCR. Mifepristone (RU-486) is a glucocorticoid and progesterone receptor antagonist. In a study conducted with a small number of patients, the efficacy of mifepristone in patients with chronic CSCR has been demonstrated. However, due to side effects such as hypotension, syncope, vaginal bleeding, and gastrointestinal disorders, it has not been a standard treatment method.

Acetazolamide

Although case reports are showing that acetazolamide treatment reduces subretinal fluid,⁵⁹ there is no increase in final BCVA and does not reduce recurrences.⁶⁰ Its use can be tried in patients who need rapid recovery.

Beta-Blocker

Case studies have reported that metoprolol, metipranolol, trimepranol, and propranolol provide improvement.⁶¹⁻⁶⁴ No significant difference was observed between the non-selective beta-blocker metoprolol and the beta-1-selective blocker metipranolol.⁶³

Rifampin

The cytochrome p450 enzyme activator rifampin is believed to treat the manifestations of CSCR by altering the metabolism of endogenous steroids. It has been shown that rifampin decreases subretinal fluid and increases visual recovery in patients with CSCR. In addition, oral rifampin is effective in reducing subretinal fluid and increasing visual recovery in patients with chronic CSCR. It is still in the hypothesis stage that rifampin accelerates the metabolism of endogenous steroids by activating the cytochrome p450 enzyme system and improves the pump function of the RPE and increases the resorption of subretinal fluid. However, the cases should be followed up in terms of gastrointestinal side effects, elevated liver enzymes, and hematological side effects while using this drug.

CONCLUSION

OCT stands out as the gold standard for imaging, diagnosis, and follow-up. Although there is no gold standard method in treatment, reduced dose and low-fluence PDT seems to be the best treatment option.

ETHICAL DECLARATIONS

Referee Evaluation Process: Externally peer-reviewed.

Conflict of Interest Statement: The authors have no conflicts of interest to declare.

Financial Disclosure: The authors declared that this study has received no financial support.

Author Contributions: All of the authors declare that they have all participated in the design, execution, and analysis of the paper, and that they have approved the final version.

REFERENCES

- Spitznas M, Huke J. Number, shape, and topography of leakage points in acute type I central serous retinopathy. *Graefes Arch Clin Exp Ophthalmol*. 1987;225(6):437-440. doi:10.1007/BF02334172
- Yamada K, Hayasaka S, Setogawa T. Fluorescein-angiographic patterns in patients with central serous chorioretinopathy at the initial visit. *Ophthalmologica*. 1992;205(2):69-76. doi:10.1159/000310315
- Bujarborua D, Nagpal PN, Deka M. Smokestack leak in central serous chorioretinopathy. *Graefes Arch Clin Exp Ophthalmol*. 2010;248(3):339-351. doi:10.1007/s00417-009-1212-5
- Yang L, Jonas JB, Wei W. Optical coherence tomography-assisted enhanced depth imaging of central serous chorioretinopathy. *Invest Ophthalmol Vis Sci*. 2013;54(7):4659-4665. Published 2013 Jul 12. doi:10.1167/iovs.12-10991
- Ellabban AA, Tsujikawa A, Ooto S, et al. Focal choroidal excavation in eyes with central serous chorioretinopathy. *Am J Ophthalmol*. 2013;156(4):673-683. doi:10.1016/j.ajo.2013.05.010
- Imamura Y, Fujiwara T, Margolis R, Spaide RF. Enhanced depth imaging optical coherence tomography of the choroid in central serous chorioretinopathy. *Retina*. 2009;29(10):1469-1473. doi:10.1097/IAE.0b013e3181be0a83
- Lehmann M, Bousquet E, Beydoun T, Behar-Cohen F. PACHYCHOROID: an inherited condition?. *Retina*. 2015;35(1):10-16. doi:10.1097/IAE.0000000000000287
- Lehmann M, Wolff B, Vasseur V, et al. Retinal and choroidal changes observed with 'En face' enhanced-depth imaging OCT in central serous chorioretinopathy. *Br J Ophthalmol*. 2013;97(9):1181-1186. doi:10.1136/bjophthalmol-2012-302974
- Piccolino FC, De La Longrais RR, Manea M, Cicinelli S. Posterior cystoid retinal degeneration in central serous chorioretinopathy. *Retina*. 2008;28(7):1008-1012. doi:10.1097/IAE.0b013e31816b4b86
- Kim HC, Cho WB, Chung H. Morphologic changes in acute central serous chorioretinopathy using spectral domain optical coherence tomography. *Korean J Ophthalmol*. 2012;26(5):347-354. doi:10.3341/kjo.2012.26.5.347
- Matsumoto H, Kishi S, Otani T, Sato T. Elongation of photoreceptor outer segment in central serous chorioretinopathy. *Am J Ophthalmol*. 2008;145(1):162-168. doi:10.1016/j.ajo.2007.08.024
- Fujimoto H, Gomi F, Wakabayashi T, Sawa M, Tsujikawa M, Tano Y. Morphologic changes in acute central serous chorioretinopathy evaluated by fourier-domain optical coherence tomography. *Ophthalmology*. 2008;115(9):1494-1500.e15002. doi:10.1016/j.ophtha.2008.01.021
- Ahlers C, Geitzenauer W, Stock G, Golbaz I, Schmidt-Erfurth U, Prunte C. Alterations of intraretinal layers in acute central serous chorioretinopathy. *Acta Ophthalmol*. 2009;87(5):511-516. doi:10.1111/j.1755-3768.2008.01468.x
- Matsumoto H, Sato T, Kishi S. Outer nuclear layer thickness at the fovea determines visual outcomes in resolved central serous chorioretinopathy. *Am J Ophthalmol*. 2009;148(1):105-10.e1. doi:10.1016/j.ajo.2009.01.018
- Iida T, Yannuzzi LA, Spaide RF, Borodoker N, Carvalho CA, Negrao S. Cystoid macular degeneration in chronic central serous chorioretinopathy. *Retina*. 2003;23(1):1-138. doi:10.1097/00006982-200302000-00001
- Landa G, Barnett JA, Garcia PM, Tai KW, Rosen RB. Quantitative and qualitative spectral domain optical coherence tomography analysis of subretinal deposits in patients with acute central serous retinopathy. *Ophthalmologica*. 2013;230(2):62-68. doi:10.1159/000350231
- Maruko I, Iida T, Ojima A, Sekiryu T. Subretinal dot-like precipitates and yellow material in central serous chorioretinopathy. *Retina*. 2011;31(4):759-765. doi:10.1097/IAE.0b013e3181fbc8e8
- von Rückmann A, Fitzke FW, Fan J, Halfyard A, Bird AC. Abnormalities of fundus autofluorescence in central serous retinopathy. *Am J Ophthalmol*. 2002;133(6):780-786. doi:10.1016/s0002-9394(02)01428-9
- Prunte C, Flammer J. Choroidal capillary and venous congestion in central serous chorioretinopathy. *Am J Ophthalmol*. 1996;121(1):26-34. doi:10.1016/s0002-9394(14)70531-8
- Iida T, Kishi S, Hagimura N, Shimizu K. Persistent and bilateral choroidal vascular abnormalities in central serous chorioretinopathy. *Retina*. 1999;19(6):508-512. doi:10.1097/00006982-199911000-00005
- Kitaya N, Nagaoka T, Hikichi T, et al. Features of abnormal choroidal circulation in central serous chorioretinopathy. *Br J Ophthalmol*. 2003;87(6):709-712. doi:10.1136/bjo.87.6.709
- Sharma T, Shah N, Rao M, et al. Visual outcome after discontinuation of corticosteroids in atypical severe central serous chorioretinopathy. *Ophthalmology*. 2004;111(9):1708-1714. doi:10.1016/j.ophtha.2004.03.025
- Yannuzzi LA. Central serous chorioretinopathy: a personal perspective. *Am J Ophthalmol*. 2010;149(3):361-363. doi:10.1016/j.ajo.2009.11.017
- Nicholson B, Noble J, Foroughian F, Meyerle C. Central serous chorioretinopathy: update on pathophysiology and treatment. *Surv Ophthalmol*. 2013;58(2):103-126. doi:10.1016/j.survophthal.2012.07.004
- Lim JW, Kang SW, Kim YT, Chung SE, Lee SW. Comparative study of patients with central serous chorioretinopathy undergoing focal laser photocoagulation or photodynamic therapy. *Br J Ophthalmol*. 2011;95(4):514-517. doi:10.1136/bjo.2010.182121
- Robertson DM, Ilstrup D. Direct, indirect, and sham laser photocoagulation in the management of central serous chorioretinopathy. *Am J Ophthalmol*. 1983;95(4):457-466. doi:10.1016/0002-9394(83)90265-9
- Maruko I, Iida T, Sugano Y, Ojima A, Ogasawara M, Spaide RF. Subfoveal choroidal thickness after treatment of central serous chorioretinopathy. *Ophthalmology*. 2010;117(9):1792-1799. doi:10.1016/j.ophtha.2010.01.023
- Peyman GA, Bok D. Peroxidase diffusion in the normal and laser-coagulated primate retina. *Invest Ophthalmol*. 1972;11(1):35-45.
- Ficker L, Vafidis G, While A, Leaver P. Long-term follow-up of a prospective trial of argon laser photocoagulation in the treatment of central serous retinopathy. *Br J Ophthalmol*. 1988;72(11):829-834. doi:10.1136/bjo.72.11.829
- Samy CN, Gragoudas ES. Laser photocoagulation treatment of central serous chorioretinopathy. *Int Ophthalmol Clin*. 1994;34(3):109-119. doi:10.1097/00004397-199403430-00012
- Robertson DM, Ilstrup D. Direct, indirect, and sham laser photocoagulation in the management of central serous chorioretinopathy. *Am J Ophthalmol*. 1983;95(4):457-466. doi:10.1016/0002-9394(83)90265-9
- Tarantola RM, Law JC, Recchia FM, Sternberg P Jr, Agarwal A. Photodynamic therapy as treatment of chronic idiopathic central serous chorioretinopathy. *Lasers Surg Med*. 2008;40(10):671-675. doi:10.1002/lsm.20720
- Chan WM, Lai TY, Lai RY, Liu DT, Lam DS. Half-dose verteporfin photodynamic therapy for acute central serous chorioretinopathy: one-year results of a randomized controlled trial. *Ophthalmology*. 2008;115(10):1756-1765. doi:10.1016/j.ophtha.2008.04.014
- Dang Y, Sun X, Xu Y, et al. Subfoveal choroidal thickness after photodynamic therapy in patients with acute idiopathic central serous chorioretinopathy. *Ther Clin Risk Manag*. 2014;10:37-43. doi:10.2147/TCRM.S54213

35. Reibaldi M, Cardascia N, Longo A, et al. Standard-fluence versus low-fluence photodynamic therapy in chronic central serous chorioretinopathy: a nonrandomized clinical trial. *Am J Ophthalmol*. 2010;149(2):307-315.e2. doi:10.1016/j.ajo.2009.08.026
36. Uetani R, Ito Y, Oiwa K, Ishikawa K, Terasaki H. Half-dose vs one-third-dose photodynamic therapy for chronic central serous chorioretinopathy. *Eye (Lond)*. 2012;26(5):640-649. doi:10.1038/eye.2012.66
37. Ober MD, Yannuzzi LA, Do DV, et al. Photodynamic therapy for focal retinal pigment epithelial leaks secondary to central serous chorioretinopathy. *Ophthalmology*. 2005;112(12):2088-2094. doi:10.1016/j.ophtha.2005.06.026
38. Yannuzzi LA, Slakter JS, Gross NE, et al. Indocyanine green angiography-guided photodynamic therapy for treatment of chronic central serous chorioretinopathy: a pilot study. *Retina*. 2003;23(3):288-298. doi:10.1097/00006982-200306000-00002
39. Inoue R, Sawa M, Tsujikawa M, Gomi F. Association between the efficacy of photodynamic therapy and indocyanine green angiography findings for central serous chorioretinopathy. *Am J Ophthalmol*. 2010;149(3):441-6.e62. doi:10.1016/j.ajo.2009.10.011
40. Lim JW, Kang SW, Kim YT, Chung SE, Lee SW. Comparative study of patients with central serous chorioretinopathy undergoing focal laser photocoagulation or photodynamic therapy. *Br J Ophthalmol*. 2011;95(4):514-517. doi:10.1136/bjo.2010.182121
41. Behnia M, Khabazkhoob M, Aliakbari S, Abadi AE, Hashemi H, Pourvahidi P. Improvement in visual acuity and contrast sensitivity in patients with central serous chorioretinopathy after macular subthreshold laser therapy. *Retina*. 2013;33(2):324-328. doi:10.1097/IAE.0b013e3182670fa3
42. Brancato R, Pratesi R, Leoni G, Trabucchi G, Vanni U. Histopathology of diode and argon laser lesions in rabbit retina. A comparative study. *Invest Ophthalmol Vis Sci*. 1989;30(7):1504-1510.
43. Lanzetta P, Furlan F, Morgante L, Veritti D, Bandello F. Nonvisible subthreshold micropulse diode laser (810 nm) treatment of central serous chorioretinopathy. A pilot study. *Eur J Ophthalmol*. 2008;18(6):934-940. doi:10.1177/112067210801800613
44. Gupta B, Elagouz M, McHugh D, Chong V, Sivaprasad S. Micropulse diode laser photocoagulation for central serous chorio-retinopathy. *Clin Exp Ophthalmol*. 2009;37(8):801-805. doi:10.1111/j.1442-9071.2009.02157.x
45. Çitirik M. Nondamaging retinal laser therapy in recurrent diabetic macular edema after anti-VEGF injections. *Turk J Med Sci*. 2021;51(5):2616-2620. doi:10.3906/sag-2007-304
46. İlhan C, Çitirik M, Özdemir M. Non-damaging retinal laser therapy in chronic central serous chorioretinopathy. *Int Ophthalmol*. 2023;10.1007/s10792-023-02691-5. doi:10.1007/s10792-023-02691-5
47. Elsner H, Pörksen E, Klatt C, et al. Selective retina therapy in patients with central serous chorioretinopathy. *Graefes Arch Clin Exp Ophthalmol*. 2006;244(12):1638-1645. doi:10.1007/s00417-006-0368-5
48. Klatt C, Elsner H, Pörksen E, et al. Selektive Retina-Therapie bei Retinopathia centralis serosa mit Pigmentepithelabhebung. *Ophthalmologe*. 2006;103(10):850-855. doi:10.1007/s00347-006-1415-7
49. Chung YR, Seo EJ, Lew HM, Lee KH. Lack of positive effect of intravitreal bevacizumab in central serous chorioretinopathy: meta-analysis and review. *Eye (Lond)*. 2013;27(12):1339-1346. doi:10.1038/eye.2013.236
50. Lim JW, Kim MU, Shin MC. Aqueous humor and plasma levels of vascular endothelial growth factor and interleukin-8 in patients with central serous chorioretinopathy. *Retina*. 2010;30(9):1465-1471. doi:10.1097/IAE.0b013e3181d8e7fe
51. Lim JW, Kim MU. The efficacy of intravitreal bevacizumab for idiopathic central serous chorioretinopathy. *Graefes Arch Clin Exp Ophthalmol*. 2011;249(7):969-974. doi:10.1007/s00417-010-1581-9
52. Chan WM, Lai TY, Liu DT, Lam DS. Intravitreal bevacizumab (avastin) for choroidal neovascularization secondary to central serous chorioretinopathy, secondary to punctate inner choroidopathy, or of idiopathic origin. *Am J Ophthalmol*. 2007;143(6):977-983. doi:10.1016/j.ajo.2007.02.039
53. Artunay O, Yuzbasioglu E, Rasier R, Sengul A, Bahcecioglu H. Intravitreal bevacizumab in treatment of idiopathic persistent central serous chorioretinopathy: a prospective, controlled clinical study. *Curr Eye Res*. 2010;35(2):91-98. doi:10.3109/02713680903428306
54. Lim JW, Ryu SJ, Shin MC. The effect of intravitreal bevacizumab in patients with acute central serous chorioretinopathy. *Korean J Ophthalmol*. 2010;24(3):155-158. doi:10.3341/kjo.2010.24.3.155
55. Wang M, Munch IC, Hasler PW, Prunte C, Larsen M. Central serous chorioretinopathy. *Acta Ophthalmol*. 2008;86(2):126-145. doi:10.1111/j.1600-0420.2007.00889.x
56. Wilkinson-Berka JL, Tan G, Jaworski K, Harbig J, Miller AG. Identification of a retinal aldosterone system and the protective effects of mineralocorticoid receptor antagonism on retinal vascular pathology. *Circ Res*. 2009;104(1):124-133. doi:10.1161/CIRCRESAHA.108.176008
57. Bousquet E, Beydoun T, Zhao M, Hassan L, Offret O, Behar-Cohen F. Mineralocorticoid receptor antagonism in the treatment of chronic central serous chorioretinopathy: a pilot study. *Retina*. 2013;33(10):2096-2102. doi:10.1097/IAE.0b013e318297a07a
58. Herold TR, Prause K, Wolf A, Mayer WJ, Ulbig MW. Spironolactone in the treatment of central serous chorioretinopathy - a case series. *Graefes Arch Clin Exp Ophthalmol*. 2014;252(12):1985-1991. doi:10.1007/s00417-014-2780-6
59. Gonzalez C. Décollements séreux rétinien. Intérêt de l'acétazolamide [Serous retinal detachment. Value of acetazolamide]. *J Fr Ophtalmol*. 1992;15(10):529-536.
60. Pikkil J, Beiran I, Ophir A, Miller B. Acetazolamide for central serous retinopathy. *Ophthalmology*. 2002;109(9):1723-1725. doi:10.1016/s0161-6420(02)01157-0
61. Avci R, Deutman AF. Die Behandlung der zentralen serösen Choroidopathie mit dem Betarezeptorenblocker Metoprolol (Vorläufige Ergebnisse) [Treatment of central serous choroidopathy with the beta receptor blocker metoprolol (preliminary results)]. *Klin Monbl Augenheilkd*. 1993;202(3):199-205. doi:10.1055/s-2008-1045583
62. Tatham A, Macfarlane A. The use of propranolol to treat central serous chorioretinopathy: an evaluation by serial OCT. *J Ocul Pharmacol Ther*. 2006;22(2):145-149. doi:10.1089/jop.2006.22.145
63. Fabianová J, Porubská M, Cepilová Z. Centrálna serózna chorioretinopatia--liečba betablokátormi [Central serous chorioretinopathy--treatment with beta blockers]. *Cesk Slov Oftalmol*. 1998;54(6):401-404.
64. Chrapek O, Spacková K, Rehák J. Léčba centrální serózní chorioretinopatie betablokátory [Treatment of central serous chorioretinopathy with beta blockers]. *Cesk Slov Oftalmol*. 2002;58(6):382-386.

Tuğçe Horozoğlu Ceran

Dr. Tuğçe Horozoğlu Ceran is graduated from Hacettepe University Faculty of Medicine (English) in 2017. She completed her residency in ophthalmology between 2018-2022 at the Health Sciences University Ulucanlar Eye Training and Research Hospital. In 2022, she received the title of FEBO with the first degree in the European Board of Ophthalmology (EBO) exam. She started to work as an ophthalmologist in the Health Sciences University Ankara Dışkapı Yıldırım Beyazıt Training and Research Hospital.

